Louisiana Ornithological Society's

LOUISIANA BIRD RECORDS COMMITTEE

Rare Bird Report Form

This form is intended as a convenience in reporting observations of species on the Louisiana Bird Records Committee (LBRC) Review List. The LBRC recommends the use of this form or a similar format when submitting records for review (to ensure that all information involving an observation is accounted for). Attach additional pages as necessary. Please print or type. Attach fieldnotes, drawings, photographs, or tape recordings, if available. Include all photos for obscurely marked species. When completed, mail to Secretary, Louisiana Bird Records Committee, c/o Museum of Natural Science, 119 Foster Hall, Louisiana State University, Baton Rouge, LA 70803-3216.

- 1. English and Scientific names: Rissa tridactyla Black-legged Kittiwake
- 3. Locality: LOUISIANA; 80 miles south of Patterson in St. Mary

 Parish in the Gulf long. 91° 56′ 15″ W, lat. 28° 38′ 48″ N

 office (specific locality)
- 4. Date(s) when observed: 16 March 1998
- 5. Time(s) of day when observed: 0740
- 6. Reporting observer and address: Daniel Christian 5720 One Perkins Place Apt. B, Baton Rouge, LA
- 7. Other observers accompanying reporter who also identified the bird(s): None
- 8. Other observers who independently identified the bird(s): None
- 9. Light conditions (position of bird in relation to shade and to direction and amount of light): The sun was in back of me. I was looking at the bird to the west
 - 10. Optical equipment (type, power, condition): Leica 10 x AZ BA in excellent condition + Bushnell Scope w/15-45 x zoom in Fair condition
 - 11. Distance to bird(s): 25 mefers when first viewed with binocular
 - 12. Duration of observation: or minutes

13.	Habitat: Ocean (Gulfof mexico)
14.	9, 3, 3, 3, 3, 3, 3, 3, 3, 3, 3, 3, 3, 3,
15.	Flying with wingbeats Fairly fast closer the a large tern. Really stood out from the surrounding sea of Herring
13.	possible, give details on total length/relative size compared to other familiar species,
	body bulk, shape, proportions, bill, eye, leg, and plumage characteristics. Stress
	features that separate it from similar species. Attach additional pages as necessary):
	Stender body and wings. Bill black, small thin, The bird tiny
	compared to nearby Herring Gulls. Eye black and ear patch
	black stood out on white head, Head Smallish. Not so noticible
	was a wide smudgy grey partial collar on neck. All black outer primeries
	+ puter prim coverts up until bend in wing where they Join with black carpal
	bar, the rest of the primaries + secondaries are white (with white tips)
~	Mantle (back) and adjoining leading edge of wing in front of black carpai
Making	bur gray. Body white Underwine white tail white except for
aclean	the tip (last Ygof retrices) black, Tail mildly forked, look's flat
Shape.	when folded. legs not seen.
5-07	(military although oxigantita
16	Voice: Did not call. RPPI Harral 14 Devesto media to be seen to be
17	. Similar species (and how are they eliminated by your description?): Bonaparte sgull has dark tip secondaries those with nave collar on neck. Little guy is black capped no collar on neck bill very sn
18	
19	. Previous experience with this species: Thousands seen off the east twest coasts a
20	. Identification aids: (list books, illustrations, other birders, etc. used in identification):
	a. at time of observation: Took notes, sketch.
	b. after observation: Seabirds" by Peter Harrison 1980
	Nat Geo. Soc. Birds of North America
21	. This description is written from:notes made during the observation;notes made
	after the observation (date:);memory.
-	Are you positive of your identification (if not then explain)?:
22	The your pooling of your mental than the property of the your pooling of the your pool
22	. Signature of reporter: Daniel Christian 17 March 1998
23	(date)
	Duration of observation April 2012