LOUISIANA ORNITHOLOGICAL SOCIETY BIRD RECORDS COMMITTEE REPORT FORM

- 1. Common and scientific name: Black-headed Grosbeak (Pheucticus melanocephalus)
- 2. Number of individuals, sexes, ages, general plumage: 1 female; unsure of age.
- 3. Locality: State (Louisiana) Parish: Orleans

Exact Location: Just outside the sewage treatment plant next to the old Crescent Acres dump.

- 4. Date(s) seen: January 11, 2004
- 5. Time(s) of day seen: 11:55 a.m.-12:05 p.m.
- 6. Reporting observer and address: Phillip Wallace

4515 S. Prieur St.

New Orleans, LA 70125-5044

- 7. Other observers accompanying reporter who also identified the bird: David Muth (Discovered the bird.)
- 8. Other observers who independently identified the bird: None that I know of.
- 9. Light conditions: The light was fair to good. There was bright sunshine, but the bird was in the shade as well as the sunlight.
- 10. Optical equipment: Zeiss 10x40 bins
- 11. Distance to the bird: 15-30 yards
- 12. Duration of observation: The bird was in view for 5-7 minutes.
- 13. Habitat (general and specific): A small wooded area along a slough just outside of the swamp adjacent to the sewage treatment plant. The woods were primarily tallow and hackberry with extensive undergrowth.
- 14. Behavior: The bird was foraging practically on the ground in lantana and as high as 25 feet in tallow and hackberry. She came in initially to an Eastern Screech-Owl tape but only after it had been playing for several minutes.

15. Description:

Size and General Impression Obviously a grosbeak of the Black-headed/Rose-breasted type.

Head The bird had a bullheaded look. It had the typical female grosbeak mask or dark brown auricular patch. The orange-buff color on the throat and upper breast continued around to the nape.

Upper Parts The upper parts were brownish with darker streaking.

Wings The wings were dark brown with two white wingbars.

Underparts The underparts were generally off-white, with very thin dark streaks on the flanks. The center of the breast was clear of streaking. The throat and upper breast were an orange-buff color.

Bill The large grosbeak-sized bill was horn colored.

- **16. Voice:** The sharp "pik" call-note was heard before the bird was seen. Muth noted it as something unusual, but Wallace wasn't sure that it wasn't just an aberrant Hairy Woodpecker call.
- 17. Similar species and how they were eliminated: The absence of streaking in the center of the breast eliminates Rose-breasted. The identification is reinforced by the orange-buff color in the throat, breast, and nape, as well as by the call note, which was different from that of the Rose-breasted, which always sounds to me like "pink."
- 18. Photographs? Muth got extensive video.
- 19. Previous experience with this species: I've seen once before in Louisiana and many times out west, as recently as summer 2003.
- 20. Aids used in identification:

 a) at time of observation: None
 b) after observation: Pyle

 21. This description written from: ______ notes made during observation; _____ x ___ notes made after observation (notes were written at 2:30 p.m. on the day of the observation before consulting any field guides.); _____ from memory.
 22. Are you positive of your identification? _____ yes___
 23. Signature of reporter: ______ Wes_____
- 24. Date and time of writing this account: February 24, 2004 5:50 p.m.

Notes written at 2:30 on Black-heald Grosbeak 1365 - 12:05 Grosberk of Nach headed / Nose-browsted Size: Orang/buff color on thoat and eight breast at very them doubt streets on flats; Breast or of streaking clear of streaking. Horn colored 5111; lang grosbeck 5)ze.
Underports of white, except as described above 5 had "poa" all wite. Muth Aust Called Stertion to the call into some the bird was seen Wallace asked it it was just a Harry Wood pecher, era Honge it dobn't south quite right for a HAWD I while wing box,