## **Louisiana Ornithological Society's**

## LOUISIANA BIRD RECORDS COMMITTEE

## Report Form

This form is intended as a convenience in reporting observations of species on the Louisiana Bird Records Committee (LBRC) Review List. The LBRC recommends the use of this form or a similar format when submitting records for review (to assure that all pertinent information is accounted for). Attach additional pages as necessary. Please print or type. Attach xerox of field notes, drawings, photographs, or tape recordings, if available. Include all photos for more obscurely marked species. When completed, mail to Secretary, Louisiana Bird Records Committee, c/o Museum of Natural Science, 119 Foster Hall, Louisiana State University, Baton Rouge, LA 70803-3216.

	and I was to have the
1.	Number of Individuals, sexes, ages, general plumage (e.g., 2 in alternate
2.	DIDMANAL PLANTS IN THE STATE OF
3.	Locality: LOUISIANA: La Salle (parish)
	Cafahoula Lake (parish)
	(specific locality)
4.	Date(s) when observed: 10-11-2006
5.	Time(s) of day when observed: ~12:00 PM
6.	Reporting observer and address: William (Bill) Vermillian
٠.	128 Saint Charles Street
(2)	Lufayette LA 70501
7.	Other observers accompanying reporter who also identified the bird(s):
8.	Other observers who independently identified the bird(s):
9.	Light conditions (position of bird in relation to shade and to direction and amount of light): Optimal, no shade, sun overhead,
10.	Optical equipment (type, power, condition):  Zeiss 10 x 40 binoculars
11.	Distance to bird(s):
	150 yards
12.	Habitat: Dredged channel (dopth 6ft?), spoilbank
13.	Habitat: Dredged channel (dopth 6H), sportoure
an	S side resetated in herbaceous veg, surround
be	Stallow water (1-6 inch dopth)

include and stress habits used in identification; relate events surrounding observation):  Flying, applored to be ferrosing; scanning  water surface; relatively slew flyth E to W
15. Description (include only what was actually seen, not what "should" have been seen; include if possibly: total length/relative size compared to other familiar species, body bulk, shape, proportions, bill, eye, leg, and plumage characteristics. Stress features that separate it from similar species): Sitting in idle airboot.  Noted gull. Upon MS pectfron through binoculars mentioned to sessionate that I had a different sull. Black wedge from primary tips to leading edge of way, while trailing edge of way, they trailing edge of way, or tighthead. No black tedge of any back, lesser coverts, por tighthead. No black tedge on their noted. Since similar to Boropath's / Franklin's an teur noted. Since similar to Boropath's / Franklin's Call. Did not planedall.  Neek white  16. Voice: NA  (appox) leftway pattern
17. Similar species (include how they were eliminated by your observation):  Block wire marking not (IM! 3 toped 05 in Block-legged left)  Leftiwolke (seem are before)  18. Photographs or tape recordings obtained? (by whom? attached?):
19. Previous experience with this species: Zeno, except for study in field guide; Hewever, that study was sufficient for me to defende that the uses the levely species before consulting me guide in identification aids: (list books, illustrations, other birders, etc. used in confirming identification):  a. at time of observation: Notional Geo. Field Gruze to the observation: National Seaboards book, Sabla produced guide, TOS Hewever of X birds, Lovery's Birds etc.  21. This description is written from:notes made during the observation (notes attached?);notes made after the observation (date:);memory.
22. Are you positive of your identification if not, explain: Yes, 23. Signature of reporter:
(date) (time)

Behavior of bird / circumstances of observation (flying, feeding, resting;

14.