See cards

1 DEC. 1998

from Parish

3

- observed with yellow-ramps
- presumed in 9
- gray back/ramp.

J. Flerence bedueen other Spp.

Jellow-rump = Yellow rump, more
Streaks on Flanks, lack
large white supercition + black
mask of 2775 pay

Hellow throat I flow throated her-

B+W warb = More streaks on Plantes, black tubide mantle lass distinct facial jurishin.

LOUISIANA ORNITHOLOGICAL SOCIETY BIRD RECORDS COMMITTEE REPORT FORM

This form is intended as a convenience in reporting observations of unusual birds. It may be used flexibly and need not be used at all. Leave blank any details not observed. Attach additional sheets if necessary. Please print or type. Attach drawings, photographys, etc.; include <u>all</u> photos if an obscurely marked species. When completed, mail to the Secretary of the LOS Bird Records Committee.

1.	Common and scientific name: Black-thronted Gray Warble (Denaroica highesters)
2.	Number of individuals, sexes, ages, general plumage (e.g. 2ad. in breeding plum-
	nga). / 6d. 9
3.	Locality: Louisiana: LA Parish: Cameron
	In small wood patch immediately be hind Johnson's Bayon High Date(s) when seen: Does to 12 1665
4.	Date(5) when seem. Detemper 11, 1115
5.	Times of day when seen: Not noted; ca. 1:00 f.m.
6.	Times of day when seen: Not noted; ca. 1:00 f.m. Reporting observer and address: Phillip walker?
	Hale S. Krieur SD., 19.0., Eli
7.	Other observers accompanying reporter who also identified the bild.
	Peter Yankey, David Muth, inter alia
8.	Peter Yankey David Muth inter alia Other observers who independently identified the bird: Larry Semo and discount
	the boat and bout Sunbu accompanied him.
9.	Light conditions (position of bird in relation to shade and to direction and amount
	of light): Adequate; overcast, internettent rain
LO.	Optical equipment (type, power, condition): Zeis, 10 x 40 5/ks
11.	Distance to bird: 20 feet
12.	Duration of observation: 10-15 minutes.
	Habitat (general and specific): Small wood patch; Chanese tollow tree

14. <u>Behavior</u> (flying, feeding, resting, etc.; include and stress any habits used in identification):

Feedby

Description (Include only what was actually seen, not what "should" have been seen.

Include if possible: total length; body bulk; shape, bill, eye, and leg characteristics; color and pattern of plumage, etc. Stress field marks):

Stics; color and pattern of plumage, etc. Stress field marks):

Yellow loves

White throat

Black streaked flooks is above and below eye; whote around eye white breast and below; hint of Cerulean type band across upper breast (indistinct and incomplete)

17.	Similar species (include how they are eliminated by your observations): Black-and-white eliminated by yellow local spot and far
	puttern
	Photographs (Was it photographed? By whom? Attached?): Vone
19.	Previous experience with this species: I've seen out West
20.	Aids: List books, illustrations, birders, etc. used in identification a) at time of observation;
	b) after observation: This description written from: notes made during observation; not made after observation; memory. Are you positive of your identification? yes . If not, explain:

Maring the state of the state of

from the same and the same and

Louisiana Ornithological Society's

LOUISIANA BIRD RECORDS COMMITTEE

Report Form

This form is intended as a convenience in reporting observations of species on the Louisiana Bird Records Committee (LBRC) Review List. The LBRC recommends the use of this form or a similar format when submitting records for review (to assure that all pertinent information is accounted for). Attach additional pages as necessary. Please print or type. Attach xerox of field notes, drawings, photographs, or tape recordings, if available. Include all photos for more obscurely marked species. When completed, mail to Secretary, Louisiana Bird Records Committee, c/o Museum of Natural Science, 119 Foster Hall, Louisiana State University, Baton Rouge, LA 70803-3216.

Jniversi	ity, Baton Rouge, LA 70803-3216.
1.	English and Scientific names: Black-throated Gray Warbler
2.	Number of Individuals, sexes, ages, general plumage (e.g., 2 in alternate plumage): / ad. 9
3.	Locality: LOUISIANA: Cameron
	Locality: LOUISIANA: Cameron (parish) (parish) (specific locality)
	(specific locality)
4.	Date(s) when observed: 17 Dec., 1995
5.	Time(s) of day when observed: 1:05-1:20
6.	Reporting observer and address: Matthew Pontiff
7.	742 Bill Churchman Loop De Ridder, LA 70634 Other observers accompanying reporter who also identified the bird(s): Gary Pentiff, Ross Carrie, Dawn Carrie, 4 at least 5 others
8.	Other observers who independently identified the bird(s): Dawn Carrie, Gary Pont. FF
9	Light conditions (position of bird in relation to shade and to direction and amount of
	light): bird was in no shade who vercast light
10.	Optical equipment (type, power, condition): Swift Ultralight 10x42, rentax PCF 8x40
11.	Distance to bird(sq:
12.	Duration of observation:
	15 minutes
134	Habitat:
•	little patch of woods filled with water

14.	Behavior of bird / circumstances of observation (flying, feeding, resting;
	include and stress habits used in identification; relate events surrounding observation):
	include and stress habits used in identification; relate events surrounding observation): bird flew twice, at other times, it mainly was feeding
•	on the Tallow Tree berries
	THE MINOW I LEE DETTIES

- 15. Description (include only what was actually seen, not what "should" have been seen; include if possibly: total length/relative size compared to other familiar species, body bulk, shape, proportions, bill, eye, leg, and plumage characteristics. Stress features bulk, shape, proportions, bill, eye, leg, and plumage characteristics. Ones house that separate it from similar species): White line through eye, dark gray crown, no central crown stripe. 2 white wingbars bray back under white line through eye, gray is darker. Gray under eye 4 small black chest band set off white throat. Black streaks on both sides of breast, not across. Yellow spot between eye + bill.
- voice: call note was higher + sharper than Yellow-rump's (heard both @ same time) 16.
- Similar species (include how they were eliminated by your observation): Bot warbler has stripe down crown, Cerulean of lacks tellow spot between eye of bill, is more blue above than gray BIK-thtd Blue warbler lacks strecked bides of is blue photographs of tape recordings obtained? (by whom? attached?): I've through eye.

 Previous experience with this species: none cheeked has yellow face, 17.

18.

- 19.
- 20. identification aids: (list books, illustrations, other birders, etc. used in identification):
 - at time of observation:
 - after observation: NG Birds of NA, Peterson west,
- Poterson East This description is written from: Unotes made during the observation 21. (<u>V</u>notes attached?); notes made after the observation (date:____); memory.
- Are you positive of your identification if not explain: 165 22.
- Signature of reporter: With World 12/18/95 7:45 p.m. (date) (time)